

Degenerated Third Order Linear Recurrences

Mario Catalani*

1 Introduction

Let us consider a third order linear recurrence

$$U_n = a_1 U_{n-1} + a_2 U_{n-2} + a_3 U_{n-3}, \quad U_0 = u_0, U_1 = u_1, U_2 = u_2.$$

Let us assume that the roots of the characteristic polynomial are real, simple, but two of them are equal in absolute value: we call this situation *degenerated*. If $\{\lambda_i\}$ are the roots let us assume that $\lambda_3 > 0$, $\lambda_1 = -\lambda_3$ and $\lambda_1 < \lambda_2 < \lambda_3$. Then the coefficients of the recurrence are $a_1 = \lambda_2$, $a_2 = \lambda_3^2$, $a_3 = -\lambda_2 \lambda_3^2$.

The closed form (generalized Binet's formula) is

$$U_n = c_1 (-\lambda_3)^n + c_2 \lambda_2^n + c_3 \lambda_3^n,$$

where $\{c_i\}$ are determined by the initial condition $\{u_i\}$, as well as by the roots $\{\lambda_i\}$.

2 Limits

Using Binet's formula the ratio of two consecutive terms is given by

$$\frac{U_n}{U_{n-1}} = \frac{c_1 (-\lambda_3)^n + c_2 \lambda_2^n + c_3 \lambda_3^n}{c_1 (-\lambda_3)^{n-1} + c_2 \lambda_2^{n-1} + c_3 \lambda_3^{n-1}}.$$

After dividing both numerator and denominator by λ_3^{n-1} we get

$$\frac{U_n}{U_{n-1}} = \frac{c_1 \lambda_3 (-1)^n + c_2 \lambda_2 \left(\frac{\lambda_2}{\lambda_3}\right)^{n-1} + c_3 \lambda_3}{c_1 (-1)^{n-1} + c_2 \left(\frac{\lambda_2}{\lambda_3}\right)^{n-1} + c_3}.$$

*Department of Economics, University of Torino, Italy. mario.catalani@unito.it

Now the limit for $n \rightarrow \infty$ depends on the parity of n . If n is even

$$L_1 = \lim_{n \rightarrow \infty} \frac{U_n}{U_{n-1}} = \lambda_3 \frac{c_1 + c_3}{-c_1 + c_3}, \quad (1)$$

while if n is odd

$$L_2 = \lim_{n \rightarrow \infty} \frac{U_n}{U_{n-1}} = \lambda_3 \frac{-c_1 + c_3}{c_1 + c_3}. \quad (2)$$

Let us express c_1 and c_3 explicitly as functions of the initial conditions. Using Binet's formula we have

$$\begin{cases} c_1 + c_2 + c_3 & = u_0, \\ -c_1\lambda_3 + c_2\lambda_2 + c_3\lambda_3 & = u_1, \\ c_1\lambda_3^2 + c_2\lambda_2^2 + c_3\lambda_3^2 & = u_2. \end{cases}$$

Let $\mathbf{A}$ be the following matrix

$$\mathbf{A} = \begin{bmatrix} 1 & 1 & 1 \\ -\lambda_3 & \lambda_2 & \lambda_3 \\ \lambda_3^2 & \lambda_2^2 & \lambda_3^2 \end{bmatrix}.$$

Let $\mathbf{c}$ the column vector with elements $\{c_i\}$ and $\mathbf{u}$ the column vector with elements $\{u_i\}$. Then

$$\mathbf{c} = \mathbf{A}^{-1} \mathbf{u}.$$

We obtain

$$\mathbf{A}^{-1} = \begin{bmatrix} \frac{\lambda_2}{2(\lambda_2 + \lambda_3)} & -\frac{1}{2\lambda_3} & \frac{1}{2\lambda_3^2 + 2\lambda_2\lambda_3} \\ \frac{\lambda_3^2}{\lambda_3^2 - \lambda_2^2} & 0 & \frac{1}{-\lambda_3^2 + \lambda_2^2} \\ \frac{\lambda_2}{-2\lambda_3 + 2\lambda_2} & \frac{1}{2\lambda_3} & \frac{1}{2\lambda_3^2 - 2\lambda_2\lambda_3} \end{bmatrix}.$$

Then

$$c_1 + c_3 = \frac{-\lambda_2^2 u_0 + u_2}{\lambda_3^2 - \lambda_2^2},$$

and

$$c_3 - c_1 = \frac{\lambda_3^2(-\lambda_2 u_0 + u_1) + \lambda_2(-\lambda_2 u_1 + u_2)}{\lambda_3^3 - \lambda_3 \lambda_2^2}.$$

It follows

$$\begin{aligned} L_1 &= \frac{\lambda_3^2(\lambda_2^2 u_0 - u_2)}{\lambda_3^2(\lambda_2 u_0 - u_1) + \lambda_2(\lambda_2 u_1 - u_2)}, \\ L_2 &= \frac{\lambda_3^2(\lambda_2 u_0 - u_1) + \lambda_2(\lambda_2 u_1 - u_2)}{\lambda_2^2 u_0 - u_2}. \end{aligned}$$

3 Consequences

If we write

$$\gamma = \frac{c_1 + c_3}{-c_1 + c_3}, \quad (3)$$

then

$$\frac{L_1}{\lambda_3} = \gamma, \quad \frac{L_2}{\lambda_3} = \frac{1}{\gamma}.$$

It follows

$$\begin{aligned} \lambda_3^2 &= L_1 L_2 \\ &= \lim_{n \rightarrow \infty} \frac{U_n}{U_{n-1}} \lim_{n \rightarrow \infty} \frac{U_{n-1}}{U_{n-2}} \\ &= \lim_{n \rightarrow \infty} \frac{U_n}{U_{n-1}} \frac{U_{n-1}}{U_{n-2}} \\ &= \lim_{n \rightarrow \infty} \frac{U_n}{U_{n-2}}. \end{aligned} \quad (4)$$

That is, this limit exists and it does not depend on the initial conditions.

Also

$$\begin{aligned} \gamma^2 &= \frac{L_1}{L_2} \\ &= \frac{\lim_{n \rightarrow \infty} \frac{U_n}{U_{n-1}}}{\lim_{n \rightarrow \infty} \frac{U_{n-1}}{U_{n-2}}} \\ &= \lim_{n \rightarrow \infty} \frac{U_n}{U_{n-1}} \frac{U_{n-2}}{U_{n-1}} \\ &= \lim_{n \rightarrow \infty} \frac{U_n U_{n-2}}{U_{n-1}^2}. \end{aligned} \quad (5)$$

That is, this limit exists and depends on the initial conditions.

4 Existence of a Limit

Under particular initial conditions our ratio converges. The equality $L_1 = L_2$ entails

$$\lambda_3^2 (\lambda_2^2 u_0 - u_2)^2 = [\lambda_3^2 (\lambda_2 u_0 - u_1) + \lambda_2 (\lambda_2 u_1 - u_2)]^2.$$

We can obtain u_2 as a function of u_0 and u_1 . This leads to a quadratic equation in u_2 whose solutions are

$$u_2 = \begin{cases} \lambda_2 \lambda_3 u_0 + (\lambda_2 - \lambda_3) u_1 \\ -\lambda_2 \lambda_3 u_0 + (\lambda_2 + \lambda_3) u_1. \end{cases}$$

If we take the first one then

$$L_1 = L_2 = -\lambda_3.$$

On the other hand if we take the second one then

$$L_1 = L_2 = \lambda_3.$$